

DNA  
DIAGNOSTIC

# HemaVision 28Q

Fast  
screening for  
Leukemia -  
associated  
translocations

## HemaVision®28Q

- Fast  
Results in just 4 hours
- Screening for 28  
translocations
- Detection of +145 break-  
points and splice variants
- CE-IVD marked test

HemaVision

Detect translocations  
in just 4 hours.

Beratung und Vertrieb in Deutschland durch:



**HiSS**  
Diagnostics

## Introduction

HemaVision®-28Q is a RT-qPCR based assay (Reverse Transcription followed by Real-Time PCR) for detection of leukemia associated fusion gene transcripts in total RNA from whole blood or bone marrow samples.

HemaVision®-28Q provides a rapid screening for 28 translocations with more than 145 clinically relevant chromosomal breakpoints. Both common and rare translocation breakpoints and alternative splice variants are detected in only 4 hours from RNA extraction.

The kit screens for rare translocations and fusion products, which cannot be detected with other commercial screening kits.

## 28 translocations

del1(p32) (STIL-TAL1)
t(1;11) (p32;q23) (KMT2A-EPS15)
t(1;11) (q21;q23) (KMT2A-MLLT11)
t(1;19) (q23;p13) (TCF3-PBX1)
t(3;5) (q25;q34) (NPM1-MLF1)
t(3;21) (q26;q22) (RUNX1-MDS1/EVI1)
t(4;11) (q21;q23) (KMT2A-AFF1)
t(5;12) (q33;p13) (ETV6-PDGFRB)
t(5;17) (q35;q21) (NPM1-RARA)
t(6;9) (p23;q34) (DEK-NUP214)
t(6;11) (q27;q23) (KMT2A-AFON)
t(8;21) (q22;q22) (RUNX1-RUNX1T1)
t(9;9) (q34;q34) (SET-NUP214)
t(9;11) (p22;q23) (KMT2A-MLLT3)
t(9;12) (q34;p13) (ETV6-ABL1)
t(9;22) (q34;q11) (BCR-ABL1)
t(10;11) (p12;q23) (KMT2A-MLLT10)
t(11;17) (q23;q21) (KMT2A-MLLT6)
t(11;17) (q23;q21) (ZBTB16-RARA)
t(11;19) (q23;p13.1) (KMT2A-ELL)
t(11;19) (q23;p13.3) (KMT2A-MLLT1)
t(12;21) (p13;q22) (ETV6-RUNX1)
t(12;22) p13;q11 (ETV6-MN1)
t(15;17) (q24;q21) (PML-RARA)
inv(16) (p13;q22) (CBFB-MYH11)
t(16;21) (p11;q22) (FUS-ERG)
t(17;19) (q22;p13) (TCF3-HLF)
t(X;11) (q13;q23) (KMT2A-FOXO4)

## How the test works?

The starting material for the HemaVision tests is total RNA extracted from whole blood or bone marrow.


RNA is added to a ready-to-use HemaVision® 28Q cDNA reaction tube and incubated for 1 hour.

cDNA template is transferred to each of the pre-mixed HemaVision® 28Q PCR reaction tubes.

Run a qPCR in a real-time PCR instrument for 2.5 hours using FAM, ROX, and Cy5 channels.

The test is completed in 4 hours after RNA extraction.

## Easy and fast workflow



1. Pipetting/2. cDNA synthesis/3. Pipetting/4.qPCR/5.Interpretation

## Catalogue numbers:

HV01-28Q

For more info visit [www.hiss-dx.de](http://www.hiss-dx.de)

or contact Tel +49 761-389 49-0 · [hiss@hiss-dx.de](mailto:hiss@hiss-dx.de)

Beratung und Vertrieb  
in Deutschland durch:



HiSS Diagnostics GmbH

Tullastr. 70  
79108 Freiburg  
Germany

Telefon: +49 761 389 49 0  
Fax: +49 761 389 49 20

Email: [hiss@hiss-dx.de](mailto:hiss@hiss-dx.de)  
Web: [www.hiss-dx.de](http://www.hiss-dx.de)